

- Walba, D. M., Richards, R. M. & Haltiwanger, R. C. *J. Am. Chem. Soc.* **104**, 3219–3221 (1982).
- Mila, F., Stafford, C. & Capponi, S. *Phys. Rev. B* **57**, 1457–1460 (1998).
- Hayashi, M. & Ebisawa, H. *J. Phys. Soc. Jpn* **70**, 3495–3498 (2001).
- Kaneda, K. & Okabe, Y. *Phys. Rev. Lett.* **86**, 2134–2137 (2001).
- Hearmon, R. F. S. *An Introduction to Applied Anisotropic Elasticity* (Oxford Univ. Press, London, 1961).
- White, J. H. *Am. J. Math.* **91**, 693–728 (1969).
- Vologodskii, A. V., Anshelevich, V. V., Lukashin, A. V. &

- Frank-Kamenetskii, M. D. *Nature* **280**, 294–298 (1979).
- Tsutsumi, K. *et al. Phys. Rev. Lett.* **39**, 1675–1676 (1977).
- Monceau, P. *Electronic Properties of Inorganic Quasi-one-dimensional Compounds* (Reidel, Dordrecht, 1985).
- Berry, M. V. *Proc. R. Soc. Lond. A* **392**, 45–57 (1984).
- Ando, T., Nakanishi, T. & Saito, R. *J. Phys. Soc. Jpn* **67**, 2857–2862 (1998).
- Iijima, S. *Nature* **354**, 56–58 (1991).
- Hyde, S. T. *Phys. Chem. Miner.* **20**, 190–200 (1993).

**Competing financial interests:** declared none.

Endosymbiotic bacteria


## GroEL buffers against deleterious mutations

**G**roEL, a heat-shock protein that acts as a molecular chaperone<sup>1</sup>, is overproduced in endosymbiotic but not in free-living bacteria<sup>2–4</sup>, presumably to assist in the folding of conformationally damaged proteins. Here we show that the overproduction of GroEL in *Escherichia coli* masks the effects of harmful mutations that have accumulated during a simulated process of vertical transmission. This molecular mechanism, which may be an adaptation to the bacterium's intracellular lifestyle, is able to rescue lineages from a progressive fitness decline resulting from the fixation of deleterious mutations under strong genetic drift<sup>5,6</sup>.

Endosymbiotic bacteria have small population sizes, do not undergo recombination, and are maternally transmitted through tight population bottlenecks<sup>7</sup>, causing the fixation of deleterious mutations due to genetic drift and hence an irreversible decline in fitness<sup>8</sup>. However, endosymbiosis is surprisingly stable and persists over long periods<sup>9</sup>, which has led to the suggestion<sup>5</sup> that *groE* (the GroEL-encoding operon) could be buffering the mutational loss of functionally active proteins because, unlike other endosymbiont genes, it is subject to strong purifying selection<sup>5</sup>.

To test this idea, we investigated the effects of overexpression of *groE* in a set of *E. coli* strains (a free-living bacterium close to several endosymbionts<sup>9</sup>) with mutations randomly accumulated throughout the genome. These spontaneous mutations were fixed by random genetic drift in a process that simulated the vertical transmission of a single endosymbiont between hosts.

We studied the accumulation of mutations in 12 replicate lines of two *E. coli* B genotypes, one of which had a 3.3-fold-increased mutation rate<sup>10</sup>. These genotypes had already adapted to a simple environment (DM25 medium) for 10,000 generations<sup>11</sup>, suggesting that mutation accumulation might result in a decline in fitness. After 3,240 generations of mutation accumulation, we measured the fitness of the strains evolved on DM25 relative to their respective ancestors. As expected, the



**Figure 1** Effect of the overexpression of the *groE* operon on the fitness of randomly mutated strains of *Escherichia coli*. **a**, The fitness values for the ancestral strains of mutators and non-mutators are the same (left); 'mutated', fitness loss after 3,240 generations of mutation accumulation; 'groE<sup>C</sup>', fitness recovery of strains in which *groE* is overexpressed. Culture conditions, fitness assays and phenotypic markers are described elsewhere<sup>11</sup>, as is the mutation-accumulation protocol<sup>12</sup>. **b**, The regulated *E. coli* *groE* operon was changed to a constitutive form (*groE<sup>C</sup>*) by putting the structural genes under the control of the  $\beta$ -lactamase gene promoter (*Pbla*). Recombinant genotypes were obtained by transduction using P1 vir (ref. 13). The *groE<sup>C</sup>* construct contains part of the flanking *yjeH* gene, the inducible tetracycline-resistance operon (*tetA/R*) for transductant selection, the *Pbla* promoter, and the *groE* coding regions (S and L). Restriction-enzyme cutting sites are indicated; P, promoter. Further details are available from the authors.

mutated strains lost fitness (Fig. 1a) as a result of the accumulation of deleterious mutations ( $\bar{W}=0.5168 \pm 0.0159$  for mutators;  $\bar{W}=0.5208 \pm 0.0087$  for non-mutators; paired *t*-tests,  $P < 0.0001$ ).

We then replaced the regulated *groE* operon of the mutated strains with a constitutive allele (*groE<sup>C</sup>*; Fig. 1b). To estimate the fitness of the *groE<sup>C</sup>* strains, we ran competition experiments against the corresponding ancestors on DM25. Surprisingly, none of the *groE<sup>C</sup>* strains reached the expected cell density during growth overnight. A simple explanation could be that an overproduction of GroEL of about  $86 \pm 16$ -fold is deleterious because it diverts amino acids away from other cellular functions.

To test this, we grew each *groE<sup>C</sup>* strain and its ancestor in DM25 supplemented with increasing concentrations of tryptone (a mixture of peptides and amino acids). We found that *groE* overexpression was deleteri-

ous in the presence of small amounts of tryptone, but that there were no significant differences in cell density between *groE<sup>C</sup>* strains and their non-mutated ancestors at tryptone concentrations of 0.1% or higher. In endosymbionts, amino acids are not limiting because these are abundant in their intracellular environment<sup>9</sup>.

To determine whether fitness estimates depend on the environment, we estimated fitness on DM25 and on DM25 + 0.1% tryptone. Both fitness estimates were correlated (partial correlation test,  $P < 0.0001$ ).

Figure 1a shows that the average fitness of the *groE<sup>C</sup>* strains derived from non-mutator strains ( $\bar{W}=0.8801 \pm 0.0214$ ) was 75.9% greater than that of the mutated strains (paired *t*-test,  $P < 0.0001$ ), but was 12% less than that of the ancestors ( $P = 0.0002$ ); the average fitness of the *groE<sup>C</sup>* strains derived from evolved mutators was  $\bar{W}=0.8152 \pm 0.0167$ , which is 61.6% greater than that of the mutated strains (paired *t*-test,  $P < 0.0001$ ) but 18.48% less than that of the ancestral strains ( $P < 0.0001$ ).

Is fitness recovery a result of the buffering of deleterious effects by GroEL, or is it simply a general benefit associated with increased concentrations of GroEL? In favour of the first possibility, the advantage of GroEL overproduction is evident only in an amino-acid-rich environment; also, if mutation compensation is occurring, we would expect a positive correlation between the extents of fitness loss and recovery, as evidenced by their partial correlation (1-tailed test,  $P = 0.0089$ ). We conclude that GroEL overexpression is likely to be of help in maintaining these endosymbionts by protecting them against the harmful effects of accumulated mutations.

**Mario A. Fares, Mario X. Ruiz-González, Andrés Moya, Santiago F. Elena, Eladio Barrio**

*Institut Cavanilles de Biodiversitat i Biologia Evolutiva and Departament de Genètica, Universitat de València, PO Box 22085, 46071 València, Spain*  
e-mail: eladio.barrio@uv.es

- Sigler, P. B. *et al. Annu. Rev. Biochem.* **67**, 581–608 (1998).
- Sato, S. & Ishikawa, H. *J. Bacteriol.* **179**, 2300–2304 (1997).
- Aksoy, S. *Insect Mol. Biol.* **4**, 32–29 (1995).
- Charles, H., Heddi, A., Guillaud, J., Nardon, C. & Nardon, P. *Biochem. Biophys. Res. Commun.* **239**, 769–774 (1997).
- Moran, N. A. *Proc. Natl Acad. Sci. USA* **93**, 2873–2878 (1996).
- Rutherford, S. L. & Lindquist, S. *Nature* **396**, 336–342 (1998).
- Funk, D. J., Wernegreen, J. J. & Moran, N. A. *Genetics* **157**, 477–489 (2001).
- Muller, H. J. *Mutat. Res.* **1**, 2–9 (1964).
- Moran, N. A. & Wernegreen, J. J. *Trends Ecol. Evol.* **15**, 321–326 (2000).
- de Visser, J. A. G. M., Zeyl, C., Gerrish, P. J., Blanchard, J. L. & Lenski, R. E. *Science* **283**, 404–406 (1999).
- Lenski, R. E., Rose, M. R., Simpson, S. C. & Tadler, S. C. *Am. Nat.* **138**, 1315–1341 (1991).
- Kibota, T. T. & Lynch, M. *Nature* **381**, 694–696 (1996).
- Zyskind, J. W. & Bernstein, S. L. *Recombinant DNA Laboratory Manual* (Academic, San Diego, 1989).

**Competing financial interests:** declared none.