

# Weakly Uniformly Rotund Banach spaces

A. Moltó, V. Montesinos, J. Orihuela and S. Troyanski \*

## Abstract

The dual space of a WUR Banach space is weakly K-analytic.

A Banach space is said to be *weakly uniformly rotund* (WUR -for short) if given sequences  $(x_n)$  and  $(y_n)$  in the unit sphere with  $\|x_n + y_n\| \rightarrow 2$  we should have  $\text{weak-lim}_n(x_n - y_n) = 0$ . This notion has become more important since Hájek proved that every WUR Banach space must be Asplund [8]. To obtain this result he uses ideas of Stegall for the equivalence between being an Asplund space and having the Radon-Nikodym property on its dual. Using this result and the Fabian-Godefroy [4] projectional resolution of the identity in the dual of an Asplund space, Fabian, Hájek and Zizler have recently showed that for a WUR Banach space  $E$  the dual space  $E^*$  is a subspace of a WCG Banach space. Indeed they proved that for a Banach space  $E$  to have an equivalent WUR norm is equivalent to the fact that the bidual unit ball  $B_{E^{**}}$ , endowed with the weak-\* topology, will be a uniform Eberlein compact [5]. Consequently they obtain that  $E$  must be LUR renormable, too [7]. The aim of this note is to provide a direct proof of the fact that every WUR Banach space  $E$  has a dual space  $E^*$  which is weakly K-analytic. This provides a topological approach to Hájek's result on the Asplundness of the space  $E$  as well as the LUR renorming consequence on  $E$  after [6].

This paper was prepared during the visit of the forth named author to the University of Valencia in the Spring term of the Academic Year 1995-96. He acknowledges his gratitude to the hospitality and facilities provided by the University of Valencia.

In this paper,  $E$  will denote a Banach space,  $E^*$  its dual,  $B_E$  its closed unit ball,  $S_E$  its unit sphere.

**Definition 1** *A Banach space  $(E, \|\cdot\|)$  is said to be uniformly Gâteaux differentiable (UGD -for short) if for every  $0 \neq x \in E$ ,*

$$\limsup_{t \rightarrow 0} \sup_{\|y\|=1} \frac{\|y + tx\| + \|y - tx\| - 2}{t} = 0.$$

---

\*The first named author has been supported in part by DGICYT Project PB91-0326, the second named author by DGICYT PB91-0326 and PB94-0535, the third named author by DGICYT PB95-1025 and DGICYT PB91-0326, the fourth named author by a grant from the "Conselleria de Cultura, Educacio i Ciencia de la Generalitat Valenciana" and by NFSR of Bulgaria Grant MM-409/94.

The following theorem is the main result of this note:

**Theorem 1** *Let  $E$  be a Banach space such that  $E^*$  has an equivalent (not necessarily dual) UGD norm (in particular, let  $E$  be WUR Banach space). Then  $E^*$  is weakly  $K$ -analytic.*

The proof is based on the following assertions

**Fact 1** (Šmul'yan, see [3, Theorem II.6.7]) *The Banach space  $E$  is WUR if and only if  $E^*$  is UGD.*

**Theorem 2** (Talagrand [9]) *Let  $K$  be a compact space. The following assertions are equivalent:*

1.  $C(K)$  is weakly  $K$ -analytic.
2. There is an increasing mapping  $\sigma \rightarrow S_\sigma$  from  $\mathbb{N}^{\mathbb{N}}$  (endowed with the product order) in the family of compact subsets of  $C(K)$  endowed with the topology of pointwise convergence, such that  $\cup\{S_\sigma : \sigma \in \mathbb{N}^{\mathbb{N}}\}$  separates points of  $K$ .

**Remark 1** In [1] the validity of the previous theorem for an arbitrary topological space is studied. In particular, for every subset  $W$  of a Banach space  $E$  it follows that  $(W, \text{weak})$  is  $K$ -analytic if and only if  $W = \cup\{S_\sigma : \sigma \in \mathbb{N}^{\mathbb{N}}\}$  and every  $S_\sigma$  is weakly compact with  $S_\sigma \subset S_\gamma$  whenever  $\sigma \leq \gamma$  in the product order. This will be the only tool necessary here from the theory of  $K$ -analytic spaces.

**Remark 2** From Theorem 1 and [6], see also [3, p. 296], we get that every WUR Banach space admits an equivalent LUR norm.

**Remark 3** From Theorem 1 it follows Hájek's [8] result asserting that every WUR Banach space is Asplund. Indeed, if we assume that  $E$  is also separable the  $K$ -analytic structure of  $(E^*, \text{weak})$  should imply that  $E^*$  is separable too. Let us explain here an easy argument following ideas from [2]: Assume  $(E^*, \text{weak})$  is  $K$ -analytic. Let  $T$  be an usco mapping from  $\mathbb{N}^{\mathbb{N}}$  into the set of subsets of  $(E^*, \text{weak})$  with  $T(\mathbb{N}^{\mathbb{N}}) = E^*$  ( $T$  can be assumed to be increasing by Remark 1). Let  $P$  be the natural projection from  $(E^*, \text{weak}^*) \times \mathbb{N}^{\mathbb{N}}$  onto  $(E^*, \text{weak})$ . Consider the restriction  $Q$  of  $P$  to  $\Sigma := \{(x, \alpha) : (x, \alpha) \in E^* \times \mathbb{N}^{\mathbb{N}}, x \in T(\alpha)\}$ . It is easy to prove that  $Q$  is continuous: let  $(x_i^*, \alpha_i)$  be a net in  $\Sigma$  such that  $(x_i^*, \alpha_i) \rightarrow (x, \alpha) \in \Sigma$ . As  $\alpha_i \rightarrow \alpha$  we can find  $\beta \in \mathbb{N}^{\mathbb{N}}$  such that  $\alpha \leq \beta$  and  $\alpha_i \leq \beta, \forall i$ . Then  $x_i \in T(\beta), x \in T(\beta)$ , and  $x_i \xrightarrow{\text{weak}^*} x$ , hence  $x_i \xrightarrow{\text{weak}} x$ . Therefore  $E^*$  is separable too. See also theorem 2.4 in [9]. With more generality, any submetrizable topological space  $X$  is analytic if and only if there is a family of compact sets  $\{S_\sigma : \sigma \in \mathbb{N}^{\mathbb{N}}\}$  in  $X$ ,  $S_\sigma \subset S_\gamma$  whenever  $\sigma \leq \gamma$  in the product order and  $X = \cup\{S_\sigma : \sigma \in \mathbb{N}^{\mathbb{N}}\}$ , [2, theorem 7].

**Proof of Theorem 1.** It is well known that  $E$  admits an equivalent WUR norm. Then  $E^*$  has an equivalent dual UGD norm. Then given  $x^* \in S_{E^*}$  and  $\epsilon > 0$ , there exists  $\delta_\epsilon(x^*) > 0$  such that

$$\|y^* + tx^*\| + \|y^* - tx^*\| \leq 2 + \epsilon|t|, \text{ if } |t| < \delta_\epsilon(x^*) \text{ and } y^* \in S_{E^*}.$$

Given a positive integer  $p$  define

$$S_p(\epsilon) := \{x^* \in S_{E^*} : \delta_\epsilon(x^*) > \frac{1}{p}\}.$$

Obviously,

$$S_1(\epsilon) \subset S_2(\epsilon) \subset \dots \subset S_p(\epsilon) \subset S_{p+1}(\epsilon) \subset \dots$$

and  $\cup_{p=1}^\infty S_p(\epsilon) = S_{E^*}$ . Let  $\alpha = (a_n) \in \mathbb{N}^{\mathbb{N}}$ . Define

$$S_\alpha := \cap_{n=1}^\infty S_{a_n}(\frac{1}{n}).$$

We have

$$S_{E^*} = \cup\{S_\alpha : \alpha \in \mathbb{N}^{\mathbb{N}}\},$$

and

$$S_\alpha \subset S_\beta, \text{ whenever } \alpha = (a_n) \leq \beta = (b_n) \text{ (i.e., } a_n \leq b_n, \forall n).$$

This sets will give us the K-analytic structure of  $E^*$  in the weak topology. Indeed, we have the following

**Claim 1** *Given  $x^{**} \in B_{E^{**}}$ ,  $\epsilon > 0$  and  $\alpha = (a_n) \in \mathbb{N}^{\mathbb{N}}$ , there is  $x \in B_E$  such that*

$$|\langle x^{**} - x, x^* \rangle| < \epsilon, \forall x^* \in S_\alpha.$$

**Proof of the claim:** Find  $n \in \mathbb{N}$  such that  $\frac{3}{n} < \epsilon$ . Pick  $y^* \in S_{E^*}$  such that

$$\langle x^{**}, y^* \rangle > 1 - \frac{1}{na_n}.$$

Find  $x \in B_E$  such that

$$\langle x, y^* \rangle > 1 - \frac{1}{na_n}.$$

Let  $x^* \in S_\alpha$ . Since  $x^* \in S_{a_n}(\frac{1}{n})$

$$\|y^* + \frac{1}{a_n}x^*\| + \|y^* - \frac{1}{a_n}x^*\| \leq 2 + \frac{1}{na_n}.$$

In particular we have

$$\langle x^{**}, y^* + \frac{1}{a_n}x^* \rangle + \langle x, y^* - \frac{1}{a_n}x^* \rangle \leq 2 + \frac{1}{na_n} \quad (1)$$

hence

$$\frac{1}{a_n} \langle x^{**} - x, x^* \rangle \leq 2 + \frac{1}{na_n} - \langle x^{**}, y^* \rangle - \langle x, y^* \rangle < \frac{3}{na_n} < \frac{\epsilon}{a_n}$$

and so

$$\langle x^{**} - x, x^* \rangle < \epsilon, \forall x^* \in S_\alpha.$$

By interchanging  $x^{**}$  and  $x$  in (1), we get

$$|\langle x^{**} - x, x^* \rangle| < \epsilon, \forall x^* \in S_\alpha.$$

and this proves the claim.

To finish the proof of the Theorem, observe that, by the Claim, each  $S_\alpha$  is weakly relatively compact since it is weak\*-relatively compact. Thus, we have

$$S_{E^*} \subset \cup \{ \overline{S_\alpha}^{\text{weak}} : \alpha \in \mathbb{N}^{\mathbb{N}} \} := W$$

and  $W$  is weakly K-analytic in  $E^*$  [Theorem 2 and Remark 1].

Consider the map

$$(W, \text{weak}) \times [0, +\infty[ \xrightarrow{\Psi} (E^*, \text{weak})$$

given by  $\Psi(x^*, t) := t.x^*$ .  $\Psi$  is continuous,  $[0, +\infty[$  is a Polish space,  $(W, \text{weak}) \times [0, +\infty[$  is K-analytic and  $\Psi(W \times [0, +\infty]) = E^*$ , so  $(E^*, \text{weak})$  is itself K-analytic. q.e.d.

## References

- [1] B. Cascales: *On K-analytic locally convex spaces*. Arch. Math. **49** (1987), 232-244.
- [2] B. Cascales and J. Orihuela *A Sequential Propety of Set-Valued Maps*. J. Mathematical Analysis and Appl. **156** (1991), 86-100.
- [3] R. Deville, G. Godefroy and V. Zizler: *Smoothness and renormings in Banach spaces*. Longman Scientific and Technical, 1993.
- [4] M. Fabian and G. Godefroy: *The dual of every Asplund admits a projectional resolution of the identity*. Studia Math. **91** (1988), 141-151.
- [5] M. Fabian, P. Hájek and V. Zizler: *On uniform Eberlein compacta and uniformly Gâteaux smooth norms*. Serdica Math. J. **23** (1997) 351-362.
- [6] M. Fabian and S. Troyanski: *A Banach space admits a locally uniformly rotund norm if its dual is a Vašák space*. Israel J. Math. **69** (1990), 214-224.
- [7] G. Godefroy, S. Troyanski, J. H. M. Whitfield and V. Zizler: *Smoothness in weakly compactly generated Banach spaces*. J. Functional Anal. **52** (1983), 344-352.
- [8] P. Hájek: *Dual renormings of Banach spaces*. Commentationes Mathematicae Universitatis Carolinae. **37** (1996), 241-253.

[9] M. Talagrand: *Espaces de Banach faiblement  $K$ -analytiques*. Annals of Mathematics, **110** (1979), 407-438.

Address: Departament d'Anàlisi Matemàtica. Universitat de València. Dr. Moliner, 50. 46100 Burjassot (València), Spain

Address: Departamento de Matemática Aplicada. E.T.S.I. Telecomunicación. Universidad Politécnica de Valencia. C/ Vera, s/n. 46071-Valencia, Spain

Address: Departamento de Análisis Matemático. Universidad de Murcia. Campus de Espinardo. Murcia, Spain.

Address: Faculty of Mathematics and Informatics. Sofia University. 5, James Bourchier blvd. 1126 Sofia, Bulgaria.